

**Dr. McCallum-MoviPrep Bowel Preparation Instructions**  
**PLEASE READ ALL OF THESE INSTRUCTIONS CAREFULLY**

At anytime should you have any questions about the instructions, please contact our office at (520) 327-3454 (during normal business hours) so a staff member can go over them with you.

**5 DAYS BEFORE YOUR EXAM**

- **STOP** taking all anticoagulants; Coumadin, Heparin, Lovenox, Plavix, Aggranox (if okay with your Primary Care Physician or Cardiologist). If you must continue them, please let us know.
- **STOP** any bulking agents; Metamucil, Citrucel, Gemfibrozil, Ginseng, Ginkgo, Garlic tablets, Vitamin E, Multi Vitamins, & Iron Supplements.

**3 DAYS BEFORE YOUR EXAM**

- **STOP EATING ALL** Nuts, Seeds, Red Meat, Leafy Vegetables, Fruit Skins, Salads, Popcorn, Beans (all kinds), Corn, Tomatoes, Salsa, Corn Chips.
- **YOU MAY HAVE:** Pasta, Eggs, White Rice, Mashed and Sweet Potatoes, Chicken, Fish, White Bread, Cheerios, Pudding, Yogurt, Milk Shakes, Cheese, Creamy Soups, Oatmeal, Bananas, Peeled Apples and Applesauce

**DAY BEFORE YOUR EXAM-Clear liquid diet only.**

- **Do not** take any oral diabetic medications the day before or the morning of your exam.
- **Insulin dependent diabetics** should take ½ the usual dose of Insulin the day before your exam and ½ the usual dose on the day of your exam. Continue to monitor blood sugar as usual.
- Take any heart and/or blood pressure medication as usual.

Drink as much of the clear liquid diet as you can throughout the day which includes; **Beverages-** Soft drinks (orange, ginger ale, cola, Sprite, 7-Up, etc.), Gatorade, Kool-Aide, strained fruit juices w/o pulp (apple, white grape, lemonade, etc.), water, tea, or coffee. **No** milk or non-dairy creamer. No beverages that are red or purple in color. Soups-low sodium chicken or beef bouillon/broth. Desserts-Hard candies, Jell-O (lemon, Lime or orange; no fruit or toppings), or popsicles (no sherbets or fruit bars).

- Lunch and Dinner- **LIQUID DIET ONLY** Drink all the clear liquids you like.
- Starting at \_\_\_\_\_ empty 1 Pouch A and 1 Pouch B into the disposable container, add lukewarm drinking water to the top line of the container. Mix to dissolve.
- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8oz), until the full liter is consumed.
- Drink 16oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep. Continue to drink clear liquids until bedtime.
- Individual responses to laxatives vary. Remain close to a toilet, as multiple bowel movements will occur.

**DAY OF YOUR EXAM-Clear liquids only as directed. NO solid food until exam has been completed.**

- Starting at \_\_\_\_\_ empty 1 Pouch A and 1 Pouch B into the disposable container, add lukewarm drinking water to the top line of the container. Mix to dissolve.
- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8oz), until the full liter is consumed.
- Drink 16oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep.
- Take any heart and/or blood pressure medications as usual.
- **STOP DRINKING CLEAR LIQUIDS 2 HOURS PRIOR TO YOUR EXAM. Nothing after \_\_\_\_\_.**
- **Please bring a list of your medications with you.**

**\*\*\*YOU MUST HAVE SOMEONE TO DRIVE YOU HOME AFTER YOUR EXAM. WE REQUEST THAT YOUR DRIVER REMAIN DURING YOUR EXAM\*\*\***

**\*\*Our facility doors open at 6:30am. Should you be scheduled for an early morning procedure, please plan your arrival accordingly.\*\***

Tucson Gastroenterology Specialists, PC  
3040 N. Swan Road, Ste. B  
Tucson, Arizona 85712  
Phone (520) 327-3454 Fax (520) 327-3431  
www.tucsongastro.com